

## Medication Use and the Complexities of Shared Care in the Treatment of DID: Opportunities and Pitfalls

### References

Brand, B. L., Classen, C. C., Lanius, R., Loewenstein, R. J., McNary, S. W., Pain, C., & Putnam, F. W. (2009). A naturalistic study of dissociative identity disorder and dissociative disorder not otherwise specified patients treated by community clinicians. *Psychological Trauma: Theory, Research, Practice, & Policy*, 1(2), 153–171.

International Society for the Study of Trauma and Dissociation. (2011). [Chu, J. A., Dell, P. F., Van der Hart, O., Cardeña, E., Barach, P. M., Somer, E., Loewenstein, R. J., Brand, B., Golston, J. C., Courtois, C. A., Bowman, E. S., Classen, C., Dorahy, M., Sar, V., Gelinas, D. J., Fine, C. G., Paulsen, S., Kluft, R. P., Dalenberg, C. J., Jacobson-Levy, M., Nijenhuis, E. R. S., Boon, S., Chefetz, R. A., Middleton, W., Ross, C. A., Howell, E., Goodwin, G., Coons, P. M., Frankel, A. S., Steele, K., Gold, S. N., Gast, U., Young, L. M., & Twombly, J.]. Guidelines for treating dissociative identity disorder in adults, third revision. *Journal of Trauma & Dissociation*, 12, 115–187.

Loewenstein, RJ (1991). Rational psychopharmacology in the treatment of multiple personality disorder. *Psychiatric Clinics of North America*, 14 (3), 721-40.

Putnam, F. W. (1989). *Diagnosis and Treatment of Multiple Personality Disorder*. New York: Guilford Press.

Lai, C.H. (2012). Dissociative identity disorder may be relieved by the combined treatments of mirtazipine and risperidone: case report of an adolescent. (letter to the editor) *J Neuropsychiatry Clin Neurosci* 24:2, E34.