

## Where to Start First? Treating Dissociative Youth with High Rate of Comorbidity

### References

- Semiz, U. B., Inanc, L., & Bezgin, C. H. (2014). Are trauma and dissociation related to treatment resistance in patients with obsessive–compulsive disorder?. *Social psychiatry and psychiatric epidemiology*, 49(8), 1287-1296.
- Putnam, F. (2013) The Role of Abusive States of Being in Interrogation, *Journal of Trauma & Dissociation*, 14:2, 147-158, DOI:10.1080/15299732.2013.724344
- Lahav, Y., Talmon, A., Ginzburg, K., & Spiegel, D. (2019). Reenacting Past Abuse–Identification with the Aggressor and Sexual Revictimization. *Journal of Trauma & Dissociation*, 1-14.
- Erik W. Baars MD , Onno van der Hart PhD , Ellert R. S. Nijenhuis PhD ,James A. Chu MD , Gerrit Glas MD PhD & Nel Draijer PhD (2010) Predicting Stabilizing TreatmentOutcomes for Complex Posttraumatic Stress Disorder and Dissociative Identity Disorder:An Expertise-Based Prognostic Model, *Journal of Trauma & Dissociation*, 12:1, 67-87, DOI:10.1080/15299732.2010.514846
- Kloet, D. V. H.-V. D., Giesbrecht, T., Wel, J. V., Bosker, W. M., Kuypers, K. P., Theunissen, E. L., ... Ramaekers, J. G. (2015). MDMA, cannabis, and cocaine produce acute dissociative symptoms. *Psychiatry Research*, 228(3), 907–912. doi: 10.1016/j.psychres.2015.04.028